

PCT / E P 9 3 / 0 1 2 1 5

The
Patent
Office

950 3612
2

The Patent Office 24 JUN 1993
Cardiff Road
Newport
Gwent
NP9 1RH

WIPO PCT

I, the undersigned, being an officer duly authorised in accordance with Section 62(3) of the Patents and Designs Act 1907, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the Patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or the inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



PRIORITY DOCUMENT

Signed

W. Russell

Dated

14th May 1993

COC1

For official .



22 MAY 1992

27MAY 92#002D4096

PAT 1 77 UC

25.00

Your reference

900-9639

9 21 09 29.7

Notes

Please type, or write in dark ink using CAPITAL letters. A prescribed fee is payable for a request for grant of a patent. For details, please contact the Patent Office (telephone 071-438 4700).

Rule 16 of the Patents Rules 1990 the main rule governing the completion and filing of this form.

Do not give trading styles, for example, 'Trading as XYZ company', nationality or former names, for example, 'formerly (known as) ABC Ltd' as these are not required.

Warning

After an application for a Patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977 and will inform the applicant if such prohibition or restriction is necessary. Applicants resident in the United Kingdom are also reminded that under Section 23, applications may not be filed abroad without written permission unless an application has been filed not less than 6 weeks previously in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction revoked.

**The
Patent
Office**

Request for grant of a Patent Form 1/77

Patents Act 1977

1 Title of invention

- 1 Please give the title of the invention **Monoclonal antibodies and their use.**

2 Applicant's details

☐ First or only applicant

- 2a If you are applying as a corporate body please give:
Corporate name **SANDOZ LTD.**

Country (and State) **Switzerland**
of incorporation, if appropriate

- 2b If you are applying as an individual or one of a partnership please give in full:

Surname

Forenames

- 2c In all cases, please give the following details:

Address **35 Lichstrasse, CH-4002 Basle,
Switzerland**

UK postcode
(if applicable)

Country **Switzerland**

ADP number **00703207001**
(if known) ✓

2d, 2e and 2f: If there are further applicants please provide details on a separate sheet of paper.

☐ **Second applicant (if any)**

2d If you are applying as a corporate body please give:
Corporate name

Country (and State
of incorporation, if
appropriate)

2e If you are applying as an individual or one of a partnership please give in full:

Surname

Forenames

2f In all cases, please give the following details:

Address

UK postcode
(if applicable)

Country

ADP number
(if known)

Ⓢ An address for service in the
United Kingdom must be supplied

Please mark correct box

Ⓢ **Address for service details**

3a Have you appointed an agent to deal with your application?

Yes ☒ No ☐ → go to 3b

↓
please give details below

Agent's name

B. A. YORKE & CO.

Agent's address

Coomb House
7 Old John's Road
Isleworth,
Middlesex TW7 6NH

Postcode

Agent's ADP
number

02981701001

1800001

3b: If you have appointed an agent, all
correspondence concerning your
application will be sent to the agent's
United Kingdom address.

3b If you have not appointed an agent please give a name and address in the
United Kingdom to which all correspondence will be sent:

Name

Address

Postcode

ADP number
(if known)

Daytime telephone
number (if available)

④ Reference number

4 Agent's or
applicant's reference
number (if applicable) 900-9639

⑤ Claiming an earlier application date

5 Are you claiming that this application be treated as having been filed on the date of filing of an earlier application?

Yes ☐ No ☒ **⇒ go to 6**

↓
please give details below

☐ number of earlier
application or patent
number

☐ filing date
(day month year)

☐ and the Section of the Patents Act 1977 under which you are claiming:

15(4) (Divisional) ☐ 8(3) ☐ 12(6) ☐ 37(4) ☐

⑥ Declaration of priority

6 If you are declaring priority from previous application(s), please give:

Country of filing	Priority application number (if known)	Filing date (day, month, year)
-------------------	---	-----------------------------------

⑥ If you are declaring priority from a
PCT Application please enter 'PCT' as
the country and enter the country
code (for example, GB) as part of the
application number.

Please give the date in all number
format, for example, 31/05/90 for
31 May 1990.

7 The answer must be "No" if:

- any applicant is not an inventor
- there is an inventor who is not an applicant, or
- any applicant is a corporate body.

8 Please supply duplicates of claim(s), abstract, description and drawing(s).

Please mark correct box(es)

9 You or your appointed agent (see Rule 90 of the Patents Rules 1990) must sign this request.

Please sign here →

A completed fee sheet should preferably accompany the fee.

7 Inventorship

7 Are you (the applicant or applicants) the sole inventor or the joint inventor?

Please mark correct box

Yes ☐

No ☒

A Statement of Inventorship on Patents Form 7/77 will need to be filed (see Rule 15).

8 Checklist

8a Please fill in the number of sheets for each of the following types of document contained in this application.

Continuation sheets for this Patents Form 1/77

no

Claim(s)

1

Description

14

Abstract

1

Drawing(s)

8

8b Which of the following documents also accompanies the application?

Priority documents (please state how many)

no

Translation(s) of Priority documents (please state how many)

no

Patents Form 7/77 – Statement of Inventorship and Right to Grant
(please state how many)

no

Patents Form 9/77 – Preliminary Examination/Search

no

Patents Form 10/77 – Request for Substantive Examination

no

9 Request

I/We request the grant of a patent on the basis of this application.

Signed

SANDOZ LTD.
Barrie Hans

Date

24/05/1992
(day month year)

Please return the completed form, attachments and duplicates where requested, together with the prescribed fee to either:

☐ The Comptroller
The Patent Office
Cardiff Road
Newport
Gwent
NP9 1RH

or

☐ The Comptroller
The Patent Office
25 Southampton Buildings
London
WC2A 1AY

Monoclonal antibodies and their use

1. Background

As the result of the introduction of the hybridoma technology monoclonal antibodies (Mabs), mostly of mouse origin, have been made to many types of human cancer. Almost most of the markers defined by xenogeneic Mab are not strictly tumor specific but are differentiation antigens shared by tumors and certain normal and/or fetal tissues. Therefore, they are best referred to as tumor associated antigens (TAAs). Whether human tumor markers detected by xenogeneic Mabs are capable of evoking an antitumor response in cancer patients, and whether such antigens are indeed related to the response to autologous tumors in cancer patients, depends on the nature of the respective TAA and is still not fully understood. TAAs which are either naturally immunogenic in the syngeneic host or can be made immunogenic might potentially be used to induce antitumor immunity for therapeutic and possibly prophylactic benefit.

One approach towards manipulating the immune system is based on idiotypic interactions. The unique antigenic determinants in and around the antigen-combining site of an Ig molecule which make one antibody distinct from another are defined as idiotopes. The totality of all idiotopes present on the variable portion of a given antibody is referred to as its idio type (id). The molecular structure of an idio type has been localized to both the complementarity determining regions and the framework regions of the variable domain and is generally but not always contributed to by both the heavy and the light chains in specific association.

Idiotypes are serologically defined entities since injection of an antibody (often referred to as Ab1) into a syngeneic, allogeneic, or xenogeneic recipient induces the production of anti-idiotypic antibodies (often referred to as Ab2). Based on the assumption that idio type/anti-idio type interactions exist physiologically a receptor-based regulation of the immune system was postulated by Niels Jerne (Ann. Immunol. 125C, 373, 1974). His network theory views the immune system as a collection of Ig molecules and receptors on T lymphocytes, each capable of recognizing an antigenic determinant (epitope) through its combining site (paratope), and each capable of being recognized by other antibodies or cell-surface receptors of the system through the idiotopes that it displays. Many studies have indeed demonstrated that idiotypic and anti-idiotypic receptors are present on the surface of both B- and T-lymphocytes as well as on secreted antibodies.

When the binding between Ab1 and Ab2 is inhibited by the antigen to which Ab1 is directed, the idio type is considered to be binding-site-related, since it involves a site on the antibody variable domain that is engaged in antigen recognition. Those idiotopes which conformationally

mimic an antigenic epitope are called the internal image of that epitope. Since both an Ab2 and an antigen bind to the relevant Ab1, they may share a similar three-dimensional conformation which represents the internal image of the given antigen.

Tumor associated antigens are often a part of "self" and evoke a very poor immune response in cancer patients. In contrast, internal image anti-idiotypic antibodies expressing three-dimensional shapes which resemble structural epitopes of the respective TAA are recognized as foreign molecules in the tumor bearing host. Therefore, the immune response raised by therapeutic or even prophylactic vaccination may cause antitumor immunity.

Monoclonal antibodies with specificity of BR55-2 (disclosed in e.g. Wistar EP 285 059, M. Blaszyk-Thurin et al., J. Biol. Chem. 262 (1987) 372-379, or Z. Steplewski et al., Hybridoma 9 (1990) 201-210) define the Lewis Y6 antigen, a carbohydrate determinant selectively expressed on a majority of human solid tumors. Based on their properties antibodies BR55-2 can be used for passive immunotherapy of, basically, epithelial cancer.

The tumor associated Lewis Y oligosaccharide determinant which is also expressed during certain stages of embryonic development is almost not immunogenic by itself. However, monoclonal anti-idiotypic antibodies (Ab 2) against BR55-2 (Ab1) with internal image properties by resembling structural epitopes of the Lewis Y antigen may be useful for induction of a protective antitumor immunity.

The present invention comprises the generation, production and characterization of murine monoclonal internal image anti-idiotypic antibodies (Ab2) to monoclonal antibodies BR55-2 (Ab1) and the use of these anti-idiotypic Mabs for active immunization against Lewis Y antigen positive cancer.

2. Generation and characterization of murine monoclonal anti-idiotypic antibodies against the Idiotypic of antibodies BR55-2

In an attempt to minimize undesired anti-isotypic immune responses, the F(ab')₂-fragment of BR55-2, murine IgG3, was chosen for immunization. For the successful generation of murine anti-id Mabs against the idiotypic of the murine Mab BR55-2, it is important to maximize the immunogenicity in order to raise an appropriate immune response in the syngeneic host. Therefore the F(ab')₂-fragment which is devoid of the Fc-part (cleavage and purification described in W 092/03165) was coupled to Keyhole Limpet Hemocyanin (KLH) as immunogenic carrier using the heterobifunctional linker N-succinimidyl-3-(2-pyridyldithio)propionate (SPDP; Pharmacia) according to described methods (J. Carlsson et al., Biochem. J. 173, 723, 1978).

Balb/c mice were immunized with this BR55-2/murine IgG3-F(ab')₂-KLH-conjugate using Freund's complete adjuvant based on a typical protocol for the generation of murine Mabs. Following repeated immunizations the murine spleen cells were fused with the murine myeloma cell line SP2/0 (for experimental details see example 1).

For an appropriate selection of the cultured hybridoma cells a sequence of tests of their supernatants was performed. This selection was based on the following criteria:

- a) Secretion rate of hybridomas by determination of the concentration of murine IgG in the supernatants (for experimental details see example 2). Cells producing high amounts of murine IgG were subcloned to single cell cultures.
- b) Binding of selected supernatants to the F(ab')₂-fragment of BR55-2/murine IgG3 (for experimental details see example 3).
- c) Inhibition of binding of BR55-2/murine IgG2a to Lewis Y antigen positive SKBR5 human breast cancer cells by selected supernatants (for experimental details see example 4).

The latter test is designed to be indicative for internal image properties of Ab2's. The murine IgG2a switch variant of BR55-2 was used for binding in order to minimize detection of Ab2's recognizing remaining constant regions of the F(ab')₂-fragment of BR55-2/murine IgG3 used for immunization. This test was performed in a quantitative manner based on the IgG concentration determined in test a) (example 2). Furthermore an excess of unspecific mouse IgG was added to this inhibition experiment in order to avoid any detection of Ab2's not specific for the idiotype of BR55-2.

Hybridomas were chosen which produce IgG with an inhibition capacity of more than 95% (inhibition of binding of BR55-2/murine IgG2a to the SKBR5 cell line).

Using the test procedures mentioned above six different hybridomas were finally selected and expanded (E4, C11, B3, B9, G6, G9). All six hybridomas produce murine IgG1 as detected by subtype ELISA using rabbit-anti-mouse IgG1/peroxidase (such as the reagent of Zymed).

All six hybridomas were cultured in roller flasks (37° C, 5% CO₂ in medium G; change of medium every 3 to 4 days) and the supernatants were collected for subsequent purification.

Each supernatant containing the respective anti-id BR55-2 Mab was purified using immunoaffinity chromatography. In general, affinity chromatography is based on the interaction between an immobilized ligand and the substance of interest. In the case of anti-idiotypic BR55-2 Mabs, the highly specific ligand for the affinity column is Mab BR55-2/murine IgG2a which binds the anti-idiotypic Mabs of choice (for experimental details see example 5).

The degree of purity of the isolated anti-id BR55-2 Mabs (E4, C11, B3, B9, G6, G9) was tested by analytical FPLC ion-exchange-chromatography, size-exclusion-chromatography, SDS-

PAGE and isoelectric focussing. Purity of all six anti-id BR55-2 Mabs was >95% (for experimental details see example 6; SDS-PAGE and isoelectric focussing are shown in figures 1 and 2).

The purified anti-id Mabs were quantitatively characterized by determination of their capacity to inhibit binding of BR55-2/murine IgG3 to the Lewis Y antigen positive SKBR5 cell line. All anti-id Mabs inhibit the binding of Ab1 to its antigen based on a 1:1 stoichiometry (for experimental details see example 7; representative results are shown in figure 3).

The final proof of the internal image properties of the anti-id BR55-2 Mabs described above and their use as surrogate for the Lewis Y carbohydrate antigen is based on their ability to generate an Ab3 response recognizing Lewis Y antigen positive cells in different species. According to the network theory of N. Jerne antibodies (Ab3) induced by immunization with internal image anti-id Mabs (Ab2) have a binding specificity similar to that of Ab1. Therefore the immune response evoked by immunization with anti-id BR55-2 Mabs should be specific for Lewis Y antigen positive tumor cells. Consequently protective antitumor immunity may be induced in man by immunization with anti-id BR55-2 Mabs.

For the investigation of the properties of an Ab3 response rabbits as well as rhesus monkeys were immunized with anti-id BR55-2 #E4 using aluminium hydroxide as adjuvant. This mild adjuvant is widely used in different vaccines for human use. As a negative control the animals were also immunized with the same amount of unspecific mouse IgG1. After four immunizations during 5 weeks sera were collected at week 9 (for experimental details see example 8). Binding of serum Ig to the Lewis Y antigen positive SKBR5 breast cancer cell line and to the Lewis Y antigen negative WM9 melanoma cell line was determined (for experimental details see examples 9 and 10).

Anti-id BR55-2 #E4 elicits a high titrated humoral immune response both in rabbits and rhesus monkeys. Serum Ig of animals immunized with anti-id BR55-2 #E4 selectively binds to the Lewis Y antigen positive tumor cell line but not to the Lewis Y antigen negative WM9 cell line. In contrast, by immunization with unspecific mouse IgG1 almost no tumor cell binding serum Ig is detected. These results are summarized in table 1. In figures 4 and 5 representative Ig-binding curves obtained with pre- and immunesera of rabbits and rhesus monkeys in a cell-ELISA (SKBR5 cells) are shown. Binding of serum Ig of animals immunized with anti-id BR55-2 #E4 to SKBR5 cells can still be detected at a serum dilution of 1:10.000.

Two years after the initial immunization course the rhesus monkeys were boosted once with anti-id BR55-2 #E4 or unspecific mouse IgG1 (for experimental details see example 8). Bind-

ing of monkey serum Ig to the Lewis Y antigen positive SKBR5 breast cancer cell line was determined before as well as 1 week and 4 weeks after boost immunization.

In rhesus monkeys treated with anti-id BR55-2 #E4 two years after first immunization Lewis Y antigen positive tumor cell binding serum Ig still can be detected. 1 week following boost immunization tremendously increased titers of serum Ig specifically binding to Lewis Y antigen positive tumor cells are found in animals boosted with anti-id BR55-2 #E4. These results are summarized in Table 2.

A similar pattern of immunoreactivity of the different monkey sera before and after first immunization course as well as before and after the boost immunization can also be demonstrated by binding to a membrane preparation of Lewis Y antigen positive SKBR5 cells (for experimental details see example 11). The results are summarized in Table 3.

In conclusion, immunization with anti-id BR55-2 Mabs leads to a high titered immune response specific for Lewis Y antigen positive cancer cells. Thereby the internal image properties of these anti-id Mabs and their use as surrogate tumor antigen for therapeutic and prophylactic active immunization with the aim of induction of antitumor immunity in man is highlighted.

The following examples illustrate the invention. The abbreviations have the following meanings:

BSA:	bovine serum albumin
CDC:	complement dependent cytotoxicity
DMEM:	Dulbecco modified Eagle Medium
ELISA:	enzyme-linked immunosorbent assay
FCS:	fetal calf serum
Mab:	monoclonal antibody
PBS:	phosphate-buffered saline
RPMI:	Rosewell Park Memorial Institute
SDS:	sodium dodecyl sulfate
KLH:	keyhole limpet hemocyanin
SPDP:	N-succinimidyl-3-(2-pyridyl-dithio-propionate)
PAGE:	polyacrylamide gel electrophoresis
IEF:	isoelectric focussing
PEG:	polyethyleneglycol

The materials referred to in the examples are as follows:

Microtiterplates: Immunoplates II (Nunc)

Cell lines: SKBR5: human breast cancer cell line
WM9: human melanoma cell line
SP2/0: mouse myeloma cell line

Medium A: RPMI 1640 + 2 g/l NaHCO_3
100 U/ml penicillin G
100 $\mu\text{g/ml}$ streptomycin sulfate
4 mM glutamine
10 % FCS (heat-inactivated, γ -globulin-free)

Medium B: RPMI 1640 + 2 g/l NaHCO_3
100 U/ml penicillin G
100 $\mu\text{g/ml}$ streptomycin sulfate
4 mM glutamine
5 % FCS (heat-inactivated)

Medium C: DMEM
10 % NCTC-135 (synthetic medium, Gibco)
1 % MEM non essential amino acids (Gibco)
0.5 % sodium pyruvate
0.5 % oxalacetic acid (Sigma)
20 % FCS (heat-inactivated)
4 mM glutamine
100 U/ml penicillin G
100 $\mu\text{g/ml}$ streptomycin sulfate

Medium D: Medium C + 1.36 mg/l hypoxanthine
0.39 mg/l thymidine

Medium E: Medium D + 0.4 mg/l aminopterin

Medium F: Medium C + mouse thymocytes (thymocytes of one Balb/c mouse resuspended in 25 ml medium C)

Medium G: DMEM
10 % FCS (heat-inactivated)
4 mM glutamine
100 U/ml penicillin G
100 $\mu\text{g/ml}$ streptomycin sulfate

PEG: polyethyleneglycol (MW = 3400)
1 g is dissolved in 1 ml DMEM

PBS deficient: 138.0 mM NaCl

1.5 mM KOH

2.7 mM KCl

6.5 mM Na_2HPO_4

pH 7.2

Coating buffer: 15 mM Na_2CO_3

35 mM NaHCO_3

3 mM NaN_3

pH 9.6

Staining buffer: 24.3 mM citric acid

51.4 mM Na_2HPO_4

pH 5.0

Washing buffer: 2% NaCl

0.2 % Triton X-100

in PBS deficient

Substrate solution: 40 mg o-phenylenediaminedihydrochloride

100 ml staining buffer

20 μl H_2O_2 30%

Binding buffer: 0.1 M Tris/HCl

0.2 M NaCl

pH 7.5

Elution buffer: 0.15 M glycine/HCl

0.2 M NaCl

pH 2.8

Coupling buffer: 0.1 M NaHCO_3

0.5 M NaCl

pH 8.0

In the following examples which illustrate the invention but in no way limit its scope references to temperature are in degrees celsius.

Example 1: Generation of anti-id BR55-2 #E4

1.1 Immunization of mice

Balb/c mice are immunized with each 100 µg of F(ab')₂-fragment of BR55-2/murine IgG3, coupled to KLH via SPDP as described (J. Carlsson et al., Biochem. J. 173, 723, 1978) by intraperitoneal injection in the following scheme:

day 0: 100 µg of conjugate (1 mg/ml in PBS def.) + 100 µl of Freund's complete adjuvant

day 7 and 28: 100 µg of conjugate (1 mg/ml in PBS def.) + 100 µl of Freund's incomplete adjuvant

On days 8, 9, 10 and 11 after primary immunization i.v. a total of 4 boost injections (each 100 µg of conjugate in 100 µl of PBS def.) are given.

On day 12 the spleens are taken out aseptically, suspended in PBS def. and washed thrice in PBS def.

1.2 Hybridization

These spleen cells are added to a suspension of SP2/O cells in a ratio 1:1 and centrifuged at 900 g for 5 minutes. 1 ml of PEG-solution (37°) is added dropwise to the cell pellet within 1 minute and diluted with 1 ml of PBS def. (37°) within the next minute. 10 ml of medium C are added under gently rotation and the suspension is diluted to 50 ml with PBS def. The suspension is centrifuged at 800 g for 5 minutes, the pellet resuspended in medium D and the cells are transferred into the wells of a microtiterplate (Nunc 96) at a concentration of 2.5×10^5 cells/well. After overnight incubation at 37°/5% CO₂ 100 µl/well of medium E are added. After 72 hours and then every four days the medium is replaced by medium D.

Example 2: Quantitative determination of mouse IgG in hybridoma supernatants

100 µl aliquots of rabbit-anti-mouse-IgG (such as the reagent of Nordic; 1:1000 in coating buffer) are added to the wells of microtiter plates, and incubated at 37° for 60 minutes.

The plates are washed 6 times with washing buffer, 200 µl of PBS def./5% FCS are added and incubated for 30 minutes at 37°. The plates are washed as described above. 100 µl aliquots of the hybridoma supernatants obtained after 2 weeks culture are added and the plates are incubated for 60 minutes at 37°. Unbound antibody is washed out as described above and 100 µl aliquots of peroxidase-conjugated antibody (rabbit-anti-mouse-IgG/peroxidase such as the reagent of Dianova; 1:1000 in PBS/2% FCS) are added.

After incubation for 30 minutes at 37° the plates are washed 4 times with washing buffer and twice with staining buffer.

100 μ l aliquots of substrate solution are added and colour development is stopped after 5 minutes with 50 μ l aliquots of 4 N H_2SO_4 /well. Photometric extinction is measured at 492 nm (reference measurement 620 nm).

Example 3: Specific binding of hybridoma supernatant-IgG to BR55-2 F(ab')₂-fragment (ELISA)

Hybridomas producing sufficient mouse IgG (i.e. more as 10-fold optical density than the medium-blank) are subcloned to single cell culture in medium F and cultured in medium G for additional 2 weeks. The supernatants are tested as follows:

100 μ l aliquots of F(ab')₂-fragment of BR55-2 (10 μ g/ml; dilution in coating buffer) are added to the wells of microtiter plates, and incubated at 37° for 60 minutes.

The plates are washed 6 times with washing buffer, 200 μ l of PBS def./5% FCS are added and incubated for 30 minutes at 37°. The plates are washed as described above. 100 μ l of hybridoma supernatants are added and the plates are incubated for 60 minutes at 37°. Unbound antibody is washed out as described above and 100 μ l aliquots of peroxidase-conjugated antibody (rabbit-anti-mouse-IgG-Fc/peroxidase such as the reagent of Dianova; 1:1000 in PBS/2 % FCS) are added.

After incubation for 30 minutes at 37° the plates are washed 4 times with washing buffer and twice with staining buffer.

100 μ l aliquots of substrate solution are added and colour development is stopped after 5 minutes with 50 μ l aliquots of 4 N H_2SO_4 /well. Photometric extinction is measured at 492 nm (reference measurement 620 nm).

Example 4: Inhibition of binding of BR55-2/murine IgG2a to SKBR5 human breast cancer cells by hybridoma supernatant-IgG (cell ELISA)

All hybridoma supernatants which are positive in the above described assay are tested as follows:

Microtiter plates are pretreated with poly-L-lysine hydrobromide (20-30 kD; 20 μ g/ml in PBS def.; 100 μ l/well; 30 minutes, room temperature), washed twice with PBS def. (200 μ l/well) and then incubated overnight at 4° with 50 μ l/well of a suspension of SKBR5 cells in medium B (4×10^6 cells/ml).

After removal of the supernatant the cells are fixed with 50 μ l of glutardialdehyde/well (0.1% in physiological saline) for 5 minutes at room temperature, the supernatants are removed, the cells resuspended in 200 μ l/well of PBS def./1% BSA/0.1% NaN_3 and left for 1 hour at room temperature.

Supernatants are removed and the plates are washed twice with 200 μ l/well of PBS containing 0.05% Tween 20. Hybridoma supernatants adjusted to 1 μ g/ml mouse IgG are preincubated with 10-fold excess of unspecific mouse IgG for 30 minutes at 37°. Then these samples are preincubated with 0.5 μ g/ml of BR55-2/murine IgG2a for 30 minutes at 37°. 100 μ l of this mixture are added to the cells and the plates are incubated for 1 hour at 37°. Unbound antibody is washed out twice with 100 μ l/well of ice-cold PBS containing 0.05 % Tween 20. 100 μ l aliquots of peroxidase-conjugated antibody (rabbit-anti-mouse IgG2a/peroxidase such as the reagent of Zymed; 1:1000 in PBS def./2% FCS) are added. After incubation for 45 minutes at 37° the wells are washed thrice with the PBS/Tween 20 solution mentioned above and then 100 μ l of the substrate solution are added to each well. After 5 minutes the colour development is stopped by addition of 50 μ l aliquots of 4 N H₂SO₄/well. Binding of the antibody to the cells is determined by measuring extinction at 492 nm (reference measurement 620 nm).

Example 5: Immunoaffinity purification of anti-Id BR55-2 Mabs

5.1 Preparation of BR55-2/murine IgG2a Sepharose

10 g of freeze dried activated CH-Sepharose 4B is suspended in 1mM HCl, transferred to a sinter glass filter and washed with 2 l of 1 mM HCl for 15 minutes. The ligand (120 mg of BR55-2/murine IgG2a) dissolved in 50 ml of coupling buffer is mixed with the washed gel in a stoppered vessel and rotated end over end for one hour at room temperature. The gel is washed with coupling buffer and incubated for one hour with 50 ml of 1M ethanolamine for blocking of any remaining active groups. The affinity sorbent is then washed with three cycles of alternating pH. Each cycle consists of a wash at pH 4 (0.1 M acetate, 0.5 M NaCl) followed by a wash at pH 8 (0.1 M Tris, 0.5 M NaCl).

5.2 Isolation of the anti-Id BR55-2 Mabs

The chromatography is performed at 4°. The column (BIO REX MP column, diameter 1.5 cm) is filled with Mab BR55-2/murine IgG2a Sepharose (volume 35 ml). The gel is washed with binding buffer and elution buffer. After equilibration with binding buffer conditioned medium containing anti-Id BR55-2 is loaded onto the column at a flow rate of 15 ml/min. After elution of the breakthrough fraction, the bound anti-Id BR55-2 is desorbed with elution buffer and neutralized immediately after desorption with 1 M Tris/HCl buffer, pH 7.5.

5.3 Concentration of the anti-Id BR55-2 Mabs

Concentration of the eluted antibody solution (0.12 mg/ml) is performed in a stirred Amicon ultrafiltration cell using a PM 10 Diaflo membrane. The solute rejection for IgG is more than 98%, the final concentration of IgG amounts to 3.7 mg/ml.

Example 6: Characterization of purified anti-Id BR55-2 Mabs

6.1 Ion-exchange-chromatography on Mono-Q

Column: Mono-Q HR5/5 (Pharmacia)
 Buffer A: 20 mM tri-ethanolamine, pH 7.7
 Buffer B: 20 mM tri-ethanolamine, 1 M NaCl, pH 7.7
 Flow rate: 1 ml/min
 Detection: UV 280 nm
 Gradient: linear 2%/min
 Results: > 95% purity found for all anti-id BR55-2 Mabs

6.2 High performance size-exclusion-chromatography

Column: Zorbax GF250, 9.4 x 250 mm
 Buffer : Sodium phosphate 0.1 M, 0.2 M NaCl, pH 7.0
 Flow rate: 1 ml/min
 Detection: UV 280 nm
 Results: > 95% purity found for all anti-id BR55-2 Mabs

6.3 SDS-PAGE

Experiments are performed both under reducing and non-reducing conditions according to the method of Laemmli using 10 % acrylamide gels (results are shown in Figure 1).

6.4 Isoelectric focussing

Analysis is performed with the Phast-system (Pharmacia) using a pH-gradient 3-9 (Phast gel IEF 3-9) and silver staining for detection of the protein bands (results are shown in figure 2).

Example 7: Binding of BR55-2/murine IgG3 to SKBR5 cell line (cell-ELISA) - Inhibition by anti-Id BR55-2 #E4

Microtiter plates are pretreated with poly-L-lysine hydrobromide (20-30 kD; 20 µg/ml in PBS def.; 100 µl/well; 30 minutes, room temperature), washed twice with PBS def. (200 µl/well) and then incubated overnight at 4° with 50 µl/well of a suspension of SKBR5 cells in medium B (4x10⁶ cells/ml). After removal of the supernatant the cells are fixed with 50 µl of glutardialdehyde/well (0.1 % in physiological saline) for 5 minutes at room temperature, the supernatants are removed, the cells resuspended in 200 µl/well of PBS def./1% BSA/0.1% NaN₃ and left for 1 hour at room temperature.

Supernatants are removed and the plates are washed twice with 200 µl/well of PBS containing 0.05 % Tween 20. Anti-id BR55-2 #E4 is diluted in PBS def. containing 2% FCS (10 to 0.5 µg/ml). To each of these dilutions 1 µg/ml of BR55-2/murine IgG3 is added. 100 µl of

this mixture are added to the cells and the plates are incubated for 1 hour at 37°. Unbound antibody is washed out twice with 100 µl/well of ice-cold PBS containing 0.05 % Tween 20. 100 µl aliquots of peroxidase-conjugated antibody (rabbit anti-mouse IgG3/peroxidase such as the reagent of Zymed; 1:1000 in PBS def./2% FCS) are added. After incubation for 45 minutes at 37° the wells are washed thrice with the PBS/Tween 20 solution mentioned above and then 100 µl of the substrate solution are added to each well. After 5 minutes colour development is stopped by addition of 50 µl of 4 N H₂SO₄/well. Binding of the antibody to the cells is determined by measuring extinction at 492 nm (reference measurement 620 nm).

Example 8: Immunization of rabbits and rhesus monkeys with anti-id BR55-2 #E4

8.1 Immunization of rabbits with anti-id BR55-2 #E4

Three female chinchilla rabbits are immunized by intradermal application of 300 µg of anti-id BR55-2 #E4 adsorbed on aluminium hydroxide (1 mg of antibody plus 3.3 mg of Al(OH)₃/ml PBS def.) on days 1, 8, 15 and 36. Three rabbits are immunized with the same amount of unspecific mouse IgG1 as negative control under the same conditions. Sera are collected before immunization and at week 9 after first immunization.

8.2 Immunization of rhesus monkeys with anti-id BR55-2 #E4

Three rhesus monkeys are immunized by subcutaneous (s.c.) application of 0.1 mg of anti-id BR55-2 #E4/kg adsorbed on aluminium hydroxide (1 mg of antibody plus 3.3 mg of Al(OH)₃/ml PBS def.) on days 1, 8, 15 and 36. Two rhesus monkeys are immunized with the same amount of unspecific mouse IgG1 as negative control under the same conditions. Sera are collected before immunization and at week 9 after first immunization.

Two years after first immunizations the same monkeys receive one s.c. boost injection with anti-id BR55-2 #E4 or unspecific mouse IgG1 respectively (same amount and formulation as above). Sera are collected before as well as 1 week and 4 weeks after boost immunization.

Example 9: Binding of rabbit serum Ig to SKBR5 cell line (cell-ELISA)

Microtiter plates are pretreated with poly-L-lysine hydrobromide (20-30 kD; 20 µg/ml in PBS def.; 100 µl/well; 30 minutes, room temperature), washed twice with PBS def. (200 µl/well) and then incubated overnight at 4° with 50 µl/well of a suspension of SKBR5 cells in medium B (4x10⁶ cells/ml). After removal of the supernatant the cells are fixed with 50 µl of glutardialdehyde/well (0.1% in physiological saline) for 5 minutes at room temperature, the supernatants are removed, the cells resuspended in 200 µl/well of PBS def./1% BSA/0.1% NaN₃ and left for 1 hour at room temperature.

Supernatants are removed and the plates are washed twice with 200 µl/well of PBS containing 0.05% Tween 20. 100 µl aliquots of rabbit sera in appropriate predilutions are added to the

cells and the plates are incubated for 1 hour at 37°. Unbound antibody is washed out twice with 100 µl/well of ice-cold PBS containing 0.05% Tween 20. 100 µl aliquots of peroxidase-conjugated antibody (goat-anti-rabbit-Ig/peroxidase such as the reagent of Dianova; 1:1000 in PBS def./2% FCS) are added. After incubation for 45 minutes at 37° the wells are washed thrice with the PBS/Tween 20 solution mentioned above and then 100 µl of the substrate solution is added to each well. After 5 minutes colour development is stopped by addition of 50 µl of 4N H₂SO₄/well. Binding of the antibody to the cells is determined by measuring extinction at 492 nm (reference measurement 620 nm).

Example 10: Binding of rhesus monkey serum Ig to SKBR5 cell line (cell-ELISA)

Microtiter plates are pretreated with poly-L-lysine hydrobromide (20-30 kD; 20 µg/ml in PBS def.; 100 µl/well; 30 minutes, room temperature), washed twice with PBS def. (200 µl/well) and then incubated overnight at 4° with 50 µl/well of a suspension of SKBR5 cells in medium B (4 x 10⁶ cells/ml). After removal of the supernatant the cells are fixed with 50 µl of glutaraldehyde/well (0.1% in physiological saline) for 5 minutes at room temperature. the supernatants are removed, the cells resuspended in 200 µl/well of PBS def./1% BSA/0.1% NaN₃ and left for 1 hour at room temperature.

Supernatants are removed and the plates are washed twice with 200 µl/well of PBS containing 0.05% Tween 20. 100 µl aliquots of rhesus monkey sera in appropriate predilutions are added to the cells and the plates are incubated for 1 hour at 37°. Unbound antibody is washed out twice with 100 µl/well of ice-cold PBS containing 0.05% Tween 20. 100 µl aliquots of peroxidase-conjugated antibody (goat-anti-human-Ig/peroxidase such as the reagent of Chemicon & Co., 1:1000 in PBS def./2% FCS) are added. After incubation for 45 minutes at 37° the wells are washed thrice with the PBS/Tween 20 solution mentioned above and then 100 µl of the substrate solution are added to each well. After 5 minutes colour development is stopped by addition of 50 µl of 4 N H₂SO₄/well. Binding of the antibody to the cells is determined by measuring extinction at 492 nm (reference measurement 620 nm).

Example 11: Binding of rhesus monkey serum Ig to a SKBR5-membrane preparation (ELISA)

Membranes of the SKBR5 human breast cancer cell line were prepared as described by D. Thom et. al., Biochem. J. 168, 187-194 (1977).

100 µl aliquots of this membrane preparation (10 µg/ml; dilution in coating buffer) are added to the wells of microtiter plates, and incubated at +4° overnight. The plates are washed 4 times with washing buffer, 200 µl of PBS def./5% FCS are added and incubated for 30 minutes at 37°. The plates are washed as described above. 100 µl aliquots of rhesus monkey sera in

appropriate dilutions in PBS def./2% FCS are added and the plates are incubated for 60 minutes at 37°. Unbound antibody is washed out as described above and 100 µl aliquots of peroxidase-conjugated antibody (goat-anti-human-Ig/peroxidase such as the reagent of Chemicon & Co., 1:1000 in PBS/2% FCS) are added. After incubation for 30 minutes at 37° the plates are washed twice with washing buffer and twice with staining buffer. 100 µl aliquots of substrate solution are added and colour development is stopped after 5 minutes with 50 µl aliquots of 4 N H₂SO₄/well. Optical density (OD) is measured at 492 nm (reference measurement 620 nm).

On view of the above experimental results anti-idiotypic BR55-2 Mabs are thus indicated for active immunization against adenocarcinomas and similar tumors, e.g. breast-, colorectal-, gastric-, pancreatic-, prostate-, ovarian- and lung cancer including small cell lung cancer, particularly for therapeutic use in humans.

For the above-mentioned use the dosage will, of course, vary depending upon e.g. the compound employed, the subject patient's age, the stage of disease, the mode of administration or the treatment desired, and can be determined by the specialist in each individual situation. It will also vary when the antibodies are used in combination with other antitumor agents or immunomodulating compounds. Administration is e.g. parenteral by subcutaneous injection possibly together with a suited adjuvant like aluminium hydroxide. The administration may be combined with immunomodulators like IL-2, CSF, INF and/or low molecular weight immunostimulants. The dosage administered is e.g. of from about 0.05 mg to about 10 mg of anti-id Mab as defined above. A typical administration schedule e.g. consists of 1 to 4 basic immunizations during 2 to 6 weeks and repeated boost immunizations e.g. every second month or on an individual basis after substantial decrease of Ab3 titers.

Claims:

1. Monoclonal murine internal image anti-idiotypic antibodies (Ab2) to monoclonal antibodies BR55-2 (Ab1).
2. Process for the production of anti-idiotypic antibodies according to claim 1 which comprises immunizing mice with BR55-2/murine IgG3-F(ab')₂-KLH-conjugate, fusing the murine spleen cells with the murine myeloma cell line SP 2/0, selecting the cultured hybridoma cells which produce IgG with an inhibition capacity of more than 95% (inhibition of binding of BR55-2 murine IgG2a to the SKBR5 cell line), purifying and isolating the anti-idiotypic antibody.
3. Use of monoclonal anti-idiotypic antibodies as defined in claim 1 for immunization against adenocarcinomas and similar tumors and against small cell lung cancer.
4. A pharmaceutical composition which comprises an anti-idiotypic monoclonal antibody as defined in claim 1 together with a pharmaceutically acceptable adjuvant, carrier or diluent for immunization against adenocarcinomas and similar tumors and against small cell lung cancer.
5. A method of immunization against adenocarcinomas and similar tumors and against small cell lung cancer, which comprises administering to a subject in need of such a treatment an effective amount of an anti-idiotypic monoclonal antibody as defined in claim 1.

Figure (c) to accompany abstract

- 16 -

900-9639 -

Abstract:

Monoclonal murine internal image anti-idiotypic antibodies (Ab2) to monoclonal antibodies BR55-2 (Ab1), process for their production and their use for immunization against adenocarcinomas and similar tumors and small cell lung cancer.

SR/hz

Table 1
Immunization of rabbits and monkeys with anti-id BR55-2 #E4 or
unspecific mouse IgG1

Relative titer increase at week 9 compared to 1:4 diluted preserum (Cell ELISA)

	Rabbit		Monkey	
	anti-id BR55-2	mouse IgG1	anti-id BR55-2	mouse IgG1
SKBR5 (Y+)	1000	16	1000	8
WM9 (Y-)	3	n.t.	25	n.t.

Table 2

Boost immunization of monkeys with anti-Id BR55-2 #E4

Binding of monkey Ig to SKBR5 cells (Cell-ELISA)

Relative titer increase at week 9 compared to 1:4 diluted preserum

9 weeks after first course	2 years after first course	1 week after boost	4 week after boost
1000	100	1500	1000

Table 3

Boost immunization of monkeys with anti-id BR55-2 #E4 or unspecific mouse IgG1

Binding of monkey Ig to a SKBR5-membrane preparation (ELISA)

Relative titer increase at week 9 compared to 1:4 diluted preserum


	9 weeks after first course	2 years after first course	1 week after boost	4 weeks after boost
anti-id BR55-2	40	1	80	50
mlgG1	1	2	n.t.	2

Figure 1
SDS - PAGE

St = Standards
EA, C11, B3, B9, G6, G9 =
anti-1d BR55-2 Mabs


St	E4	C11	B3	B9	G6	G9	St	E4	C11	B3	B9	G6	G9
reducing conditions						non-reducing conditions							

Figure 2
Isoelectric Focussing




St = Standard
R = Reference Mab
E4, C11, B3, B9, G6 and G9 = anti-id BR55-2 Mabs

Figure 3
Binding of BR55-2/murine IgG3 to SKBR5 cell line
Inhibition by anti-Id BR55-2 #E4




BR 55-2/murine IgG3: 1 μ g/ml

Figure 4
Binding of rabbit Ig to SKBR5 before and after immunization with
anti-id BR55-2 #E4 (Cell-ELISA)



—+— before immunization —o— week 9 after immun.

Figure 5
Binding of rhesus monkey Ig to SKBR5 before and after immunization with
anti-Id BR55-2 #E4 (Cell-ELISA)



—+— before immunization —x— week 9 after immun.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.